

***INVESTIGATIONS OF HYGROSCOPIC GROWTH AND PHASE
TRANSITIONS OF ATMOSPHERIC PARTICLES BY NONCONTACT
ATOMIC FORCE MICROSCOPY (AFM)***

Ocko, B., Oatis, S., Strasberg, M., Schwartz, S. and Checco, A.

Presented at the
American Physical Society March Meeting,
New Orleans, LA
March 10-14, 2008

Environmental Sciences Department/Atmospheric Sciences Division
Brookhaven National Laboratory
P.O. Box, Upton, NY
www.bnl.gov

ABSTRACT

Aerosol particles (nanometer to micrometer sized particles suspended in air) affect atmospheric radiation and cloud microphysics. A correct description of their behavior in the atmosphere is essential to accurate climate modeling. The processes by which initially hydrophobic particles become hygroscopic, accrete water from the vapor, undergo phase transition from solid particles to solution droplets are important but not well understood at a fundamental level. We have carried out AFM studies to measure changes in particle size and morphology as a function of the relative humidity for particles of sodium chloride (a substance whose bulk hygroscopic properties are well characterized) deposited on substrates with differing surface energies (Silicon Oxide, Carboxy- and Methyl-terminated organic thin-films). For particles with height >50 nm, deliquescence was observed with a relative humidity near $75 \pm 2\%$, is consistent with measurements of suspended aerosols. These preliminary results demonstrate that environmental AFM is a viable probe for studying the hygroscopic behavior of salt nanoparticles on solid supports.

NOTICE: This manuscript has been authored by employees of Brookhaven Science Associates, LLC under Contract No. DE-AC02-98CH10886 with the U.S. Department of Energy. The publisher by accepting the manuscript for publication acknowledges that the United States Government retains a non-exclusive, paid-up, irrevocable, world-wide license to publish or reproduce the published form of this manuscript, or allow others to do so, for United States Government purposes.